

Tribimaximal Neutrino Mixing and a Relation Between Neutrino- and Charged Lepton-Mass Spectra

Yoshio Koide

Department of Physics, University of Shizuoka, 52-1 Yada, Shizuoka 422-8526, Japan

E-mail address: koide@u-shizuoka-ken.ac.jp

Abstract

Brannen has recently pointed out that the observed charged lepton masses satisfy the relation $m_e + m_\mu + m_\tau = \frac{2}{3}(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2$, while the observed neutrino masses satisfy the relation $m_{\nu 1} + m_{\nu 2} + m_{\nu 3} = \frac{2}{3}(-\sqrt{m_{\nu 1}} + \sqrt{m_{\nu 2}} + \sqrt{m_{\nu 3}})^2$. It is discussed what neutrino Yukawa interaction form is favorable if we take the fact pointed out by Brannen seriously.

1 Introduction

It is well-known that the observed charged lepton mass spectrum [1] satisfies the relation [2, 3]

$$m_e + m_\mu + m_\tau = \frac{2}{3} (\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2, \quad (1.1)$$

with remarkable precision, while the observed masses (m_{f1}, m_{f2}, m_{f3}) of the other fundamental particles f_i (quarks and neutrinos) do not satisfy [4] a relation similar to (1.1) straightforwardly. However, Brannen [5] has recently pointed out a possibility that the observed neutrino masses can satisfy the relation

$$m_{\nu 1} + m_{\nu 2} + m_{\nu 3} = \frac{2}{3} (-\sqrt{m_{\nu 1}} + \sqrt{m_{\nu 2}} + \sqrt{m_{\nu 3}})^2. \quad (1.2)$$

Of course, we cannot extract the values of the neutrino mass ratios $m_{\nu 1}/m_{\nu 2}$ and $m_{\nu 2}/m_{\nu 3}$ from the neutrino oscillation data Δm_{solar}^2 and Δm_{atm}^2 unless we have more information on the neutrino masses, so that we cannot judge whether the observed neutrino masses satisfy the relation (1.2) or not. Nevertheless, it is worthwhile to examine the Brannen's speculation seriously, because it seems to bring a new view into a lepton mass matrix model if his conjecture is correct.

The Brannen's conjecture (1.2) is somewhat unforeseen, because since we have known the existence of the lepton flavor mixing, we expect that the neutrino mass matrix structure will be quite different from that of the charged leptons, so that the neutrino mass spectrum will also be different from that of the charged leptons. If the charged lepton masses satisfy the relation (1.1), the neutrino masses will not be able to satisfy the relation (1.2) even if we allow the replacement $\sqrt{m_{\nu i}} \rightarrow -\sqrt{m_{\nu i}}$. In order to understand the formula (1.1), for example, a model [3, 6, 7] with a seesaw-type mass generation mechanism [8] has been proposed:

$$M_f = m_L M_F^{-1} m_R, \quad (1.3)$$

where M_F are hypothetical heavy fermion mass matrices and, for the charged lepton sector, the structure $M_E \propto \text{diag}(1, 1, 1)$ is assumed. (For neutrino sector, Eq. (1.3) reads $M_\nu = m_L M_N^{-1} m_L^T$.) Here, the mass matrices m_L and m_R are given by $m_R \propto m_L \propto \text{diag}(v_1, v_2, v_3)$, where v_i are vacuum expectation values (VEVs) of 3-family scalars ϕ_{Li} (ϕ_{Ri}) and the VEVs v_i satisfy the relation

$$v_1^2 + v_2^2 + v_3^2 = \frac{2}{3} (v_1 + v_2 + v_3)^2. \quad (1.4)$$

Of course, we do not rule out a possibility that the values of v_i are negative. (Such a Higgs potential model which gives the relation (1.4) is proposed in Ref.[3, 7, 9, 10].) In such a seesaw-type model with diagonal m_L and m_R , mixings among fermions f_i are caused by the non-diagonal structure M_F . If M_E in the charged lepton sector is proportional to a unit matrix $\mathbf{1}$, the observed neutrino mixings tell us that the structure of the heavy neutrino matrix (the Majorana mass matrix of the right-handed neutrinos N_R) M_N cannot be unit matrix, so that the eigenvalues of the matrix $m_L M_N^{-1} m_L^T$ will not satisfy the relation (1.2) even if we allow the replacement $\sqrt{m_{\nu i}} \rightarrow -\sqrt{m_{\nu i}}$.

If we take the Brannen's speculation seriously, we must abandon such a model (1.3) (with a universal m_L structure). For example, we may modify the model as

$$M_f = m_L^f M_F^{-1} m_R^f, \quad (1.5)$$

where M_F have a unit matrix structure universally, at least, for the charged lepton and neutrino sectors, and m_L^f (and m_R^f) have flavor-dependent structures. Then, in order to give the relations (1.1) and (1.2), it is required that the eigenvalues v_{fi} of the matrices m_L^f (m_R^f) always satisfy the relation (1.4). We relax the constraint on m_L from a rigid ansatz of the universal structure into the constraint (1.4).

Since $M_F \propto \mathbf{1}$ do not play any essential role in the modified model (1.5), we may rather consider a Frogatt-Nielsen [11] type model with six dimensional operators $\bar{f}_{Li} \phi_{fi} H_L \phi_{fi} f_{Ri}$, where H_L is the conventional $\text{SU}(2)_L$ -doublet Higgs scalar, and ϕ_{fi} are 3-family $\text{SU}(2)_L$ -singlet scalars. However, in the present paper, we do not have interest in whether the effective mass matrix form originates in a seesaw model or in a Frogatt-Nielsen model. Our interest is only in the neutrino Yukawa interaction form. For convenience, in the present paper, we will discuss a possible model in the framework of a seesaw mass matrix model.

The purpose of the present paper is not to give a model which leads to the Brannen's conjecture. The purpose is to investigate what Yukawa interaction form is required when we accept the Brannen's conjecture and when we take the observed neutrino mixing into consideration. In the present paper, it is assumed that the eigenvalues v_{fi} of the Dirac mass matrices m_L^f of the charged leptons and neutrinos satisfy the relation (1.4). In the next section, we will introduce a useful parametrization for describing charged lepton and neutrino masses in terms of a permutation symmetry S_3 [12]. In Sec. 3, we will investigate a possible neutrino mass spectrum within the framework of the S_3 symmetry. In Sec. 4, we will propose a phenomenological neutrino Yukawa coupling form, where the S_3 symmetry is explicitly broken. Finally, Sec. 5 is devoted in the concluding remarks.

2 Mass spectrum parameters z_i

It is convenient to use parameters z_i which are defined by $\langle \phi_{Li}^0 \rangle \equiv v_i = vz_i$ with the normalization condition $z_1^2 + z_2^2 + z_3^2 = 1$. The relation (1.1) [(1.4)] requires that the parameters z_i satisfy the relation

$$z_1^2 + z_2^2 + z_3^2 = \frac{2}{3}(z_1 + z_2 + z_3)^2. \quad (2.1)$$

The values of z_i are obtained from the observed charged lepton masses [1] as

$$\frac{z_1}{\sqrt{m_e}} = \frac{z_2}{\sqrt{m_\mu}} = \frac{z_3}{\sqrt{m_\tau}} = \frac{1}{\sqrt{m_e + m_\mu + m_\tau}}, \quad (2.2)$$

and the explicit numerical values are $z_1 = 0.016473$, $z_2 = 0.236869$ and $z_3 = 0.971402$.

Since the relation (2.1) is invariant under any exchange $z_i \leftrightarrow z_j$, it is useful to use the language of the permutation symmetry S_3 : We define the singlet ϕ_σ and doublet (ϕ_π, ϕ_η) of S_3 for the fields ϕ_i (and also for f_i):

$$\phi_\pi = \frac{1}{\sqrt{2}}(\phi_3 - \phi_2), \quad \phi_\eta = \frac{1}{\sqrt{6}}(2\phi_1 - \phi_2 - \phi_3), \quad \phi_\sigma = \frac{1}{\sqrt{3}}(\phi_3 + \phi_2 + \phi_1), \quad (2.3)$$

where we have taken the basis (e_1, e_2, e_3) as $(e_1, e_2, e_3) = (e, \mu, \tau)$ corresponding to the definition (2.2). The definition (2.3) is different from the conventional one. This definition will become useful for description of the neutrino mixing as we show later [for example, see Eq. (3.3)].

In order that the VEVs v_i satisfy the relation (1.4), it needs that the Higgs potential leads to the relation

$$z_\pi^2 + z_\eta^2 = z_\sigma^2, \quad (2.4)$$

where $\langle \phi_\pi^0 \rangle = vz_\pi$, $\langle \phi_\eta^0 \rangle = vz_\eta$ and $\langle \phi_\sigma^0 \rangle = vz_\sigma$. The explicit form of the Higgs potential which leads to the relation (2.4) has been given in Ref.[9, 10].

By the way, in general, the eigenvalues λ_i of any Hermitian matrix M ($MM^\dagger$ if M is not Hermitian) can be expressed by the following form:

$$\begin{aligned} \lambda_1 &= \frac{1}{3}a - \frac{1}{3}b \sin \theta, \\ \lambda_2 &= \frac{1}{3}a - \frac{1}{3}b \sin \left(\theta + \frac{2}{3}\pi \right), \\ \lambda_3 &= \frac{1}{3}a - \frac{1}{3}b \sin \left(\theta + \frac{4}{3}\pi \right), \end{aligned} \quad (2.5)$$

where

$$a = \text{Tr}M, \quad b = \sqrt{2} \sqrt{3\text{Tr}(M^2) - (\text{Tr}M)^2}. \quad (2.6)$$

Therefore, any three-flavor mass spectrum can always be expressed by the two parameters b/a and θ independently of the structure of M . Only for the case with $\text{Tr}(M^2) = \frac{2}{3}(\text{Tr}M)^2$, it gives the relation $b = \sqrt{2}a$. Inversely, if we take $b = \sqrt{2}a$ in (2.5), the matrix M satisfies the relation

$\text{Tr}(M^2) = \frac{2}{3}(\text{Tr}M)^2$. (Although the author [13] and Brannen [5] have discussed a specific mass matrix form

$$\frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + \frac{1}{\sqrt{6}} \begin{pmatrix} 0 & e^{i\theta} & e^{-i\theta} \\ e^{-i\theta} & 0 & e^{i\theta} \\ e^{-i\theta} & e^{i\theta} & 0 \end{pmatrix}, \quad (2.7)$$

in connection with the mass formula (1.1), the constraint $b = \sqrt{2}a$ has been assumed (not derived) in their models, so that it does not mean that the relation (1.1) was derived in their models.)

In the present paper, since we take the three-flavor Higgs potential model [10] which gives the relation (2.1) [i.e. (2.4)], the parameters z_i can be expressed as

$$\begin{aligned} z_1 &= \frac{1}{\sqrt{6}} - \frac{1}{\sqrt{3}} \sin \theta_e, \\ z_2 &= \frac{1}{\sqrt{6}} - \frac{1}{\sqrt{3}} \sin \left(\theta_e + \frac{2}{3}\pi \right), \\ z_3 &= \frac{1}{\sqrt{6}} - \frac{1}{\sqrt{3}} \sin \left(\theta_e + \frac{4}{3}\pi \right), \end{aligned} \quad (2.8)$$

where the numerical value of θ_e is given by

$$\theta_e = \frac{\pi}{4} - \varepsilon = 42.7324^\circ \quad (\varepsilon = 2.2676^\circ), \quad (2.9)$$

from the observed charged lepton masses.

Since we assume that the neutrino masses $m_{\nu i}$ satisfy the relation (1.2), we can also define the z_i^ν parameters in the neutrino sector as

$$\begin{aligned} z_1^\nu &= \frac{1}{\sqrt{6}} - \frac{1}{\sqrt{3}} \sin \theta_\nu, \\ z_2^\nu &= \frac{1}{\sqrt{6}} - \frac{1}{\sqrt{3}} \sin \left(\theta_\nu + \frac{2}{3}\pi \right), \\ z_3^\nu &= \frac{1}{\sqrt{6}} - \frac{1}{\sqrt{3}} \sin \left(\theta_\nu + \frac{4}{3}\pi \right), \end{aligned} \quad (2.10)$$

with $|z_1^\nu| < |z_2^\nu| < |z_3^\nu|$. Then, Brannen [5] has also empirically found that if the observed neutrino mass values are given by

$$\theta_\nu = \theta_e + \frac{\pi}{12} = 57.7324^\circ, \quad (2.11)$$

which gives

$$z_1^\nu = -0.079938, \quad z_2^\nu = 0.385404, \quad z_3^\nu = 0.9192788. \quad (2.12)$$

The Brannen's empirical relation (2.11) [i.e. the z_i^ν -values (2.12)] predicts

$$R \equiv \frac{|m_{\nu 2}^2 - m_{\nu 1}^2|}{|m_{\nu 3}^2 - m_{\nu 2}^2|} = \frac{(z_2^\nu)^4 - (z_1^\nu)^4}{(z_3^\nu)^4 - (z_2^\nu)^4} = 0.0318, \quad (2.13)$$

which is in good agreement with the observed value of R [14, 15]

$$R = \frac{\Delta m_{solar}^2}{\Delta m_{atm}^2} = \frac{(7.9_{-0.5}^{+0.6}) \times 10^{-5}}{(2.72_{-0.25}^{+0.38}) \times 10^{-3}} = (2.9 \pm 0.5) \times 10^{-2}. \quad (2.14)$$

However, note that we cannot predict the ratio R only under the assumption (1.2), i.e. unless we also assume the value of θ_ν . In Fig. 1, we illustrate the predicted value of R versus θ_ν under the assumption (1.2). The present observed values of Δm_{ij}^2 are consistent with the value of $\theta_\nu \simeq 55^\circ - 58^\circ$. In other words, if we want to consider a model in which the neutrino masses satisfy the relation (1.2), we must build a model which gives $\theta_\nu \simeq 55^\circ - 58^\circ$.

In the present paper, we take the Brannen's conjecture (2.11) seriously, and we investigate what Yukawa interaction from in the neutrino sector can lead to the relation (2.11). Brannen[5] has tried to build a new mass model with an algebraic approach. However, in the present paper, we will discuss a possible mass matrix form within the framework of a conventional mass matrix model, i.e. based on a Higgs mechanism and an extended seesaw mechanism.

3 S_3 symmetry and Yukawa interaction in the neutrino sector

We have assumed that the Yukawa interaction in the charged lepton sector is given by

$$H_e = y_e \left(\bar{\ell}_{L1} E_{R1} \phi_{L1}^d + \bar{\ell}_{L2} E_{R2} \phi_{L2}^d + \bar{\ell}_{L3} E_{R3} \phi_{L3}^d \right), \quad (3.1)$$

with a universal coupling constant y_e , where $\ell_{Li} = (\nu_{Li}, e_{Li})$ and $\phi_{Li}^d = (\phi_{Li}^0, \phi_{Li}^-)$. (In the previous section, the neutrino ν_i ($i = 1, 2, 3$) denoted the mass eigenstates. However, in the present section, we will use the same notation ν_i as the $SU(2)_L$ partners of $e_{Li} = (e_L, \mu_L, \tau_L)$, respectively.) We also assume the same structure for $\bar{L}_{Ri} e_{Ri}^c \phi_{Ri}$, where $L_{Ri} = (E_{Ri}^c, N_{Ri}^c)$ and $\phi_{Ri}^d = (\phi_{Ri}^+, \bar{\phi}_{Ri}^0)$. The explicit quantum number assignments are found, for example, in Refs.[7, 16] for $SU(2)_L \times SU(2)_R \times U(1)$ model, and in Ref.[17] for $SO(10) \times SO(10)$ model.

The Yukawa interaction (3.1) is invariant under the S_3 symmetry, but, of course, it is not a general form which is invariant under the S_3 symmetry. Only when we consider that the Yukawa interaction (3.1) with the VEV $\langle \phi_{Li} \rangle = v_i = v z_i$ gives the mass matrix m_L^e in the seesaw model (1.5) with $M = M_0 \mathbf{1}$, the predicted charged lepton masses satisfy the formula (1.1).

The present observed neutrino data strongly suggest that the neutrino mixing is almost described by the so-called tribimaximal mixing [18]

$$U = \begin{pmatrix} \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{pmatrix}. \quad (3.2)$$

Under the definition (2.3), the neutrino states $(\nu_e, \nu_\mu, \nu_\tau)$ are represented by

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \begin{pmatrix} \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} \nu_\eta \\ \nu_\sigma \\ \nu_\pi \end{pmatrix}. \quad (3.3)$$

Therefore, it is useful to express the neutrino (Dirac) matrix M_ν^D on the basis $(\nu_\eta, \nu_\sigma, \nu_\pi)$, not on the basis $(\nu_\pi, \nu_\eta, \nu_\sigma)$. Here, the neutrino states ν_π , ν_η and ν_σ are defined by a relation similar

to Eq.(2.3) with $(\nu_1, \nu_2, \nu_3) = (\nu_e, \nu_\mu, \nu_\tau)$. When the mass matrix M_ν becomes diagonal on the basis $(\nu_\eta, \nu_\sigma, \nu_\pi)$, the mixing matrix is given by the tribimaximal mixing (3.2) exactly.

If we require the S_3 symmetry, the Yukawa interaction is generally given by the form

$$H_\nu = y_1 \frac{\bar{\ell}_\pi N_\pi + \bar{\ell}_\eta N_\eta + \bar{\ell}_\sigma N_\sigma}{\sqrt{3}} \phi_\sigma^u + y_2 \left[\frac{\bar{\ell}_\pi N_\eta + \bar{\ell}_\eta N_\pi}{\sqrt{2}} \phi_\pi^u + \frac{\bar{\ell}_\pi N_\pi - \bar{\ell}_\eta N_\eta}{\sqrt{2}} \phi_\eta^u \right] \\ + y_3 \left[\frac{\bar{\ell}_\pi \phi_\pi^u + \bar{\ell}_\eta \phi_\eta^u}{\sqrt{2}} N_\sigma + \bar{\ell}_\sigma \frac{\phi_\pi^u N_\pi + \phi_\eta^u N_\eta}{\sqrt{2}} \right] + y_4 \frac{\bar{\ell}_\pi N_\pi + \bar{\ell}_\eta N_\eta - 2\bar{\ell}_\sigma N_\sigma}{\sqrt{6}} \phi_\sigma^u, \quad (3.4)$$

where the heavy neutrinos N_i denote ν_{Ri} , and $\phi^u = (\bar{\phi}_L^+, \bar{\phi}_L^0)$. In the present investigation, the S_3 symmetry is a necessary condition, but it is not a sufficient condition. For the charged lepton sector, we have assumed the form (3.1), which corresponds to the case with $y_1 = \sqrt{2}y_2 = y_3 = y_e/\sqrt{3}$ and $y_4 = 0$ in the S_3 invariant general form (3.4) with $\bar{\ell}_a N_a \rightarrow \bar{\ell}_a e_{Ra}$ and $\phi_a^u \rightarrow \phi_a^d$ ($a = \pi, \eta, \sigma$). Therefore, in order to obtain the form (3.1), we must a further selection rule in addition to the S_3 invariance, for example, a cyclic permutation symmetry, and so on. For the neutrino sector, in order to reduce the number of parameters, we assume the following selection rules:

(i) The VEVs of the up-type Higgs scalars ϕ_i^u satisfy the relation

$$\langle \phi_\pi^u \rangle^2 + \langle \phi_\eta^u \rangle^2 = \langle \phi_\sigma^u \rangle^2, \quad (3.5)$$

as well as Eq.(2.4). (We assume a similar Higgs potential structure for the up-type Higgs scalars ϕ_i^u as well as ϕ_i^d .)

(ii) We have assumed the universality of the Yukawa coupling constants for the (e_1, e_2, e_3) basis in the charged lepton sector as shown in Eq.(3.1). We also assume the universality of the Yukawa coupling constants in the neutrino sector [however, not for the (ν_1, ν_2, ν_3) basis, but for the $(\nu_\pi, \nu_\eta, \nu_\sigma)$ basis].

Considering that the observed mixing is almost given by the tribimaximal mixing (3.2), i.e. the neutrino Dirac mass matrix M_ν^D is almost diagonal on the $(\nu_\eta, \nu_\sigma, \nu_\pi)$ basis, we confine ourselves to investigating a specific case with $y_1 = y_2 = y_\nu$ and $y_3 = y_4 = 0$:

$$H_\nu = y_\nu \frac{\bar{\ell}_\pi N_\pi + \bar{\ell}_\eta N_\eta + \bar{\ell}_\sigma N_\sigma}{\sqrt{3}} \phi_\sigma^u + y_\nu \left[\frac{\bar{\ell}_\pi N_\eta + \bar{\ell}_\eta N_\pi}{\sqrt{2}} \phi_\pi^u + \frac{\bar{\ell}_\pi N_\pi - \bar{\ell}_\eta N_\eta}{\sqrt{2}} \phi_\eta^u \right]. \quad (3.6)$$

The interaction (3.6) yields the neutrino Dirac mass matrix M_ν^D

$$M_\nu^D = m_0^\nu \begin{pmatrix} \frac{z_\pi^u}{\sqrt{3}} - \frac{z_\eta^u}{\sqrt{2}} & 0 & \frac{z_\pi^u}{\sqrt{2}} \\ 0 & \frac{z_\pi^u}{\sqrt{3}} & 0 \\ \frac{z_\pi^u}{\sqrt{2}} & 0 & \frac{z_\pi^u}{\sqrt{3}} + \frac{z_\eta^u}{\sqrt{2}} \end{pmatrix}, \quad (3.7)$$

where $m_0^\nu = y_\nu v_u$, $\langle \phi_\pi^u \rangle = v_u z_\pi^u$, $\langle \phi_\eta^u \rangle = v_u z_\eta^u$, $\langle \phi_\sigma^u \rangle = v_u z_\sigma^u$, and $(z_\pi^u)^2 + (z_\eta^u)^2 + (z_\sigma^u)^2 = 1$. The mass eigenvalues are given by

$$\begin{aligned} (m_\eta^\nu)' &= \left(\frac{1}{\sqrt{3}} z_\sigma^u - \frac{1}{\sqrt{2}} \sqrt{(z_\pi^u)^2 + (z_\eta^u)^2} \right) m_0^\nu, \\ m_\sigma^\nu &= \frac{1}{\sqrt{3}} z_\sigma^u m_0^\nu, \\ (m_\pi^\nu)' &= \left(\frac{1}{\sqrt{3}} z_\sigma^u + \frac{1}{\sqrt{2}} \sqrt{(z_\pi^u)^2 + (z_\eta^u)^2} \right) m_0^\nu. \end{aligned} \quad (3.8)$$

When we use the relation (3.5), i.e.

$$(z_\pi^u)^2 + (z_\eta^u)^2 = (z_\sigma^u)^2 = \frac{1}{2}, \quad (3.9)$$

we obtain

$$\begin{aligned} (m_\eta^\nu)' &= \left(\frac{1}{\sqrt{6}} - \frac{1}{2} \right) m_0^\nu, \\ m_\sigma^\nu &= \frac{1}{\sqrt{6}} m_0^\nu, \\ (m_\pi^\nu)' &= \left(\frac{1}{\sqrt{6}} + \frac{1}{2} \right) m_0^\nu, \end{aligned} \quad (3.10)$$

independently of the parameter value of z_π^u/z_η^u (in other words, independently of the magnitude of the $\nu_\pi \leftrightarrow \nu_\eta$ mixing). The result (3.10) means

$$\theta_\nu = \frac{\pi}{3}, \quad (3.11)$$

from the definition of θ_ν , Eq.(2.11).

As seen in Fig. 1, although the present interaction form could not give the Brannen's relation (2.11), the value $\theta_\nu = 60^\circ$ is very close to the Brannen's conjecture $\theta_\nu = 57.73^\circ$. In fact, the result (3.10) predicts

$$R = \frac{m_{\nu 2}^2 - m_{\nu 1}^2}{m_{\nu 3}^2 - m_{\nu 2}^2} = \frac{4\sqrt{6} - 9}{4\sqrt{6} + 9} = 0.04245. \quad (3.12)$$

Although the value (3.12) is somewhat large comparing with the Brannen's prediction (2.14), the prediction(3.11) is, at present, not ruled out within three sigma.

The value $\langle \phi_\pi^u \rangle \neq 0$ yields the ν_π - ν_η mixing:

$$\tan 2\theta_{\pi\eta} = \frac{z_\pi^u}{z_\eta^u}. \quad (3.13)$$

If we take $z_\pi^u/z_\eta^u = z_\pi^d/z_\eta^d$, the case yields a large deviation from the tribimaximal mixing. Therefore, we cannot choose the same values z_i^u as z_i^d . We must consider $\langle \phi_\pi^u \rangle \simeq 0$ differently from the case of the down-type Higgs scalars ϕ_i^d .

4 Phenomenological neutrino Yukawa interaction form

In the previous section, we have required the S_3 invariance for the neutrino Yukawa interaction H_ν , and, in order to obtain the result (3.11) near to the Brannen's relation (2.11), we have found that we need a VEV structure $\langle \phi^u \rangle \simeq 0$ differently from the VEV structure of ϕ^d . In the present section, we assume the same structure of $\langle \phi_i^u \rangle$ as $\langle \phi_i^d \rangle$ (i.e. the same parameter values of z_i). Alternatively, we abandon the S_3 invariance of H_ν , although we still use the notation (π, η, σ) in the S_3 symmetry. The purpose of the present section is not to derive the Brannen's relation (2.11) from a model with some symmetry, but to investigate what Yukawa interaction form is required if we take the Brannen's relation (2.11) and the tribimaximal mixing (3.2) seriously. Of course, if we introduce several adjustable parameters, it is always possible. Therefore, we still adhere to the universality of the coupling constants.

We consider only terms which keep the mass matrix diagonal on the $(\nu_\eta, \nu_\sigma, \nu_\pi)$ basis:

$$\begin{aligned}
H_\nu = & y_\nu \frac{\bar{\ell}_\pi N_\pi + \bar{\ell}_\eta N_\eta + \bar{\ell}_\sigma N_\sigma}{\sqrt{3}} \phi_\sigma + y_\nu \frac{\bar{\ell}_\pi N_\pi - \bar{\ell}_\eta N_\eta}{\sqrt{2}} \phi_x \\
& + y_\nu \frac{\bar{\ell}_\pi N_\pi + \bar{\ell}_\eta N_\eta - 2\bar{\ell}_\sigma N_\sigma}{\sqrt{6}} \phi_y.
\end{aligned} \tag{4.1}$$

Here, we have assumed that the first term in (4.1) is still invariant under the S_3 symmetry. The three $(\bar{\ell}N)$ terms in (4.1) are linearly independent of each other. Similarly, we assume that the scalars ϕ_x and ϕ_y are linearly independent of ϕ_σ , i.e. ϕ_x and ϕ_y are given by linearly independent combinations of ϕ_π and ϕ_η .

In order to fix ϕ_x and ϕ_y , we assume that the interaction H_ν is invariant under the exchange $\nu_\pi \leftrightarrow \nu_\eta$ ($\phi_\pi \leftrightarrow \phi_\eta$), i.e. we assume

$$\phi_x = \frac{\phi_\pi - \phi_\eta}{\sqrt{2}}, \quad \phi_y = \frac{\phi_\pi + \phi_\eta}{\sqrt{2}}. \tag{4.2}$$

Although this symmetry is analogous to the so-called $2 \leftrightarrow 3$ flavor symmetry [19] in the neutrino mass matrix, the present π - η symmetry does not mean $\nu_\mu \leftrightarrow \nu_\tau$ symmetry.

As a result, we obtain

$$\begin{aligned}
H_\nu = & y_\nu \bar{\ell}_\pi N_\pi \left[\frac{1}{\sqrt{3}} \phi_\sigma - \sqrt{\frac{2}{3}} \left(-\frac{\sqrt{3}+1}{2\sqrt{2}} \phi_\pi + \frac{\sqrt{3}-1}{2\sqrt{2}} \phi_\eta \right) \right] \\
& + y_\nu \bar{\ell}_\eta N_\eta \left[\frac{1}{\sqrt{3}} \phi_\sigma - \sqrt{\frac{2}{3}} \left(\frac{\sqrt{3}-1}{2\sqrt{2}} \phi_\pi - \frac{\sqrt{3}+1}{2\sqrt{2}} \phi_\eta \right) \right] \\
& + y_\nu \bar{\ell}_\sigma N_\sigma \left[\frac{1}{\sqrt{3}} \phi_\sigma - \sqrt{\frac{2}{3}} \left(\frac{1}{\sqrt{2}} \phi_\pi + \frac{1}{\sqrt{2}} \phi_\eta \right) \right].
\end{aligned} \tag{4.3}$$

Since we obtain

$$\langle \phi_\pi \rangle = vz_\pi = \frac{v}{\sqrt{2}} \cos \theta_e, \quad \langle \phi_\eta \rangle = vz_\eta = -\frac{v}{\sqrt{2}} \sin \theta_e, \quad \langle \phi_\sigma \rangle = vz_\sigma = \frac{v}{\sqrt{2}}, \quad (4.4)$$

from the definitions (2.3) and (2.8), the neutrino Yukawa interactions (4.3) yields the following neutrino Dirac mass matrix M_ν^D on the $(\nu_\eta, \nu_\sigma, \nu_\pi)$ basis:

$$M_\nu^D = \text{diag}(m_{\eta\eta}, m_{\sigma\sigma}, m_{\pi\pi}), \quad (4.5)$$

$$\begin{aligned} m_{\eta\eta} &= y_\nu v \left[\frac{1}{\sqrt{6}} - \frac{1}{\sqrt{3}} \sin \left(\theta_e + \frac{\pi}{12} \right) \right], \\ m_{\sigma\sigma} &= y_\nu v \left[\frac{1}{\sqrt{6}} + \frac{1}{\sqrt{3}} \sin \left(\theta_e - \frac{\pi}{4} \right) \right], \\ m_{\pi\pi} &= y_\nu v \left[\frac{1}{\sqrt{6}} + \frac{1}{\sqrt{3}} \cos \left(\theta_e - \frac{\pi}{12} \right) \right], \end{aligned} \quad (4.6)$$

where we have used that $\sin(\pi/12) = (\sqrt{3}-1)/2\sqrt{2}$ and $\cos(\pi/12) = (\sqrt{3}+1)/2\sqrt{2}$. Comparing the expression (4.6) with the expression (2.10), we obtain the Brannen's empirical relation (2.11), $\theta_\nu = \theta_e + \pi/12$, which can give a reasonable prediction (2.13). Note that the definition of the (π, η, σ) basis of S_3 , (2.3), is essential for the derivation of the Brannen's relation (2.11). If we took another conventions of the S_3 representation, we would not obtain the Brannen's relation (2.11).

If we regard $m_{\nu 3}$ as $m_{\nu 3} = \sqrt{\Delta m_{atm}^2}$, from $m_{\nu i} \propto (z_i^\nu)^2$, we obtain

$$m_{\nu 1} = (3.94_{-0.43}^{+0.27}) \times 10^{-4} \text{ eV}, \quad m_{\nu 2} = (9.17_{-0.43}^{+0.62}) \times 10^{-3} \text{ eV}, \quad m_{\nu 3} = (5.22_{-0.25}^{+0.35}) \times 10^{-2} \text{ eV}, \quad (4.7)$$

where we have used the best fit value [15] $\Delta m_{atm}^2 = (2.72_{-0.25}^{+0.38}) \times 10^{-3} \text{ eV}^2$. The predicted values (4.7) lead to

$$\Delta m_{21}^2 = (8.39_{-0.77}^{+1.17}) \times 10^{-5} \text{ eV}^2, \quad (4.8)$$

which is in good agreement with the observed value [14] $\Delta m_{21}^2 = (7.9_{-0.5}^{+0.6}) \times 10^{-5} \text{ eV}^2$.

Of course, since the mass matrix M_ν^D is diagonal in the $(\nu_\eta, \nu_\sigma, \nu_\pi)$ basis, the neutrino mixing matrix U is exactly given by the tribimaximal mixing (3.2), which gives

$$\sin^2 2\theta_{23} = 1, \quad (4.9)$$

$$\tan^2 \theta_{12} = \frac{1}{2} \quad (\theta_{12} = 35.26^\circ), \quad (4.10)$$

$$|U_{13}| = 0. \quad (4.11)$$

The prediction (4.10) is also in good agreement with the observed value [14] $\tan^2 \theta_{21} = 0.45_{-0.07}^{+0.09}$. We would like to emphasize that, in order to obtain the tribimaximal mixing (3.2), the magnitudes of the mass eigenvalues $|m_{\eta\eta}| < |m_{\sigma\sigma}| < |m_{\pi\pi}|$ are essential together with the definition of the S_3 basis (2.3).

5 Concluding remarks

In the present paper, it has been pointed out that if we take the phenomenological relations (1.2) and (2.11) seriously, we must consider that the Yukawa interactions in the neutrino sector

are given by the form (4.1). The Yukawa interactions (4.1) leads to the tribimaximal mixing (3.2) and the Brannen's relation (2.11) [so that the neutrino mass spectrum (4.7)]. Although, in Sec.4, we have put a requirement of the $\pi \leftrightarrow \eta$ symmetry for the neutrino Yukawa interaction H_ν instead of the S_3 symmetry, the requirement is, of course, merely a phenomenological assumption. For the charged lepton sector, we have assumed the S_3 invariant Yukawa interaction, while, for the neutrino sector, we have assumed the S_3 -breaking interaction (4.1). The tribimaximal mixing (3.2) has already been derived, for example, from an A_4 symmetry [20]. The interaction form (4.1) will be understood from an extended finite symmetry.

To the contrary, if we adhere the S_3 symmetry, as we discussed in Sec.3, we can obtain the result (3.11), which is numerically near to the Brannen's relation (2.11). However, we must assume that $\langle \phi_\pi^u \rangle \simeq 0$ in the up-type Higgs scalars differently from the case in the down-type Higgs scalars. The case (3.6) is also interesting.

In the present seesaw model, the 3-family $SU(2)_L$ -doublet scalars ϕ_i cause a flavor-changing neutral current (FCNC) problem. Besides, the new heavy fermions E_{Ri} and E_{Li} together with the scalars ϕ_i considerably affect the evolution of the gauge coupling constants. If we want to avoid these problems, we can take an alternative model, a Frogatt-Nielsen-like [11] model:

$$H_{eff} = y_e \bar{\ell}_L H_L^d \left(\frac{\phi}{\Lambda} \right)^2 e_R + y_\nu \bar{\ell}_L H_L^u \frac{\phi}{\Lambda} \nu_R + \bar{\nu}_R M_R \nu_R^*, \quad (5.1)$$

where $\ell_L = (\nu_L, e_L)$, $H_L^d = (H_d^+, H_d^0)$, $H_L^u = (H_u^0, H_u^-)$, ϕ is a 3-family $SU(2)_L$ -singlet scalar, and Λ is a scale of the effective theory. (Here, we have denoted the expression (5.1) symbolically. For example, the interaction $\bar{\ell}\phi\nu_R$ should read the interaction (4.1).)

In the present paper, we have discussed the masses and mixings only in the lepton sectors. We think that if there is a beautiful law in the masses and mixings of the fundamental fermions, we will find it just in the lepton sectors, because the mass generation mechanism seems to be simple just in the lepton sectors. If we define

$$R_f(\eta_1, \eta_2) = \frac{\frac{2}{3} (\eta_1 \sqrt{m_{f1}} + \eta_2 \sqrt{m_{f2}} + \sqrt{m_{f3}})^2}{m_{f1} + m_{f2} + m_{f3}}, \quad (5.2)$$

where $\eta_i = \pm 1$ ($i = 1, 2$), the ratio b/a in the expression (2.5) with $m_{fi} \propto \lambda_i^2$ is given by

$$\frac{b}{\sqrt{2}a} = \sqrt{\frac{2-R}{R}}, \quad (5.3)$$

where R (not R_f) is defined by Eq.(2.13) and a and b are defined by Eq.(2.5) [or Eq.(2.6)]. In the lepton sectors, we have found

$$R_e(+, +) = R_\nu(-, +) = 1, \quad (5.4)$$

so that $(b/\sqrt{2}a)_e = (b/\sqrt{2}a)_\nu = 1$. However, as seen in Table 1, in the quark sectors, there is no solution of a/b which gives $R_u(\eta_1^u, \eta_2^u) = R_d(\eta_1^d, \eta_2^d)$. Therefore, the present idea in the lepton

sectors cannot be applied to the quark sectors straightforwardly. In the seesaw model (1.5), we have assumed that the heavy fermion mass matrices M_F in the lepton sectors are structureless, i.e. $M \propto \mathbf{1}$. For quark sectors, we may consider that M_F have some structure except for the unit matrix $\mathbf{1}$, so that the quark masses will not satisfy the relation (5.2) with $R_f = 1$. Our goal is to find a unified description of the quark and lepton masses and mixings. For this purpose, the present study in the lepton sectors will provide a promising clue to the unified model.

Acknowledgements

The author would like to thank C. Brannen for communicating his preliminary results prior to publication. This work is supported in part by the Grant-in-Aid for Scientific Research, Ministry of Education, Science and Culture, Japan (No.18540284).

References

- [1] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. **B592**, 1 (2004).
- [2] Y. Koide, Lett. Nuovo Cimento **34**, 201 (1982); Phys. Lett. **B120**, 161 (1983); Phys. Rev. **D28**, 252 (1983).
- [3] Y. Koide, Mod. Phys. Lett. **A5**, 2319 (1990).
- [4] N. Li and B. Q. Ma, Phys. Lett. **B609**, 309 (2005); Phys. Rev. **D73**, 013009 (2006); Z. Z. Xing and H. Zhang, Phys. Lett. **B635**, 107 (2006).
- [5] C. Brannen, <http://brannenworks.com/MASSES2.pdf>, (2006).
- [6] Y. Koide and H. Fusaoka, Z. Phys. **C71**, 459 (1996).
- [7] Y. Koide and M. Tanimoto, Z. Phys. **C72**, 333 (1996).
- [8] The seesaw mechanism for charged particles is known as the “universal seesaw mechanism”: Z. G. Berezhiani, Phys. Lett. **129B**, 99 (1983); Phys. Lett. **150B**, 177 (1985); D. Chang and R. N. Mohapatra, Phys. Rev. Lett. **58**, 1600 (1987); A. Davidson and K. C. Wali, Phys. Rev. Lett. **59**, 393 (1987); S. Rajpoot, Mod. Phys. Lett. **A2**, 307 (1987); Phys. Lett. **191B**, 122 (1987); Phys. Rev. **D36**, 1479 (1987); K. S. Babu and R. N. Mohapatra, Phys. Rev. Lett. **62**, 1079 (1989); Phys. Rev. **D41**, 1286 (1990); S. Ranfone, Phys. Rev. **D42**, 3819 (1990); A. Davidson, S. Ranfone and K. C. Wali, Phys. Rev. **D41**, 208 (1990); I. Sogami and T. Shinohara, Prog. Theor. Phys. **86**, 1031 (1991); Phys. Rev. **D47**, 2905 (1993); Z. G. Berezhiani and R. Rattazzi, Phys. Lett. **B279**, 124 (1992); P. Cho, Phys. Rev. **D48**, 5331 (1993); A. Davidson, L. Michel, M. L. Sage and K. C. Wali, Phys. Rev. **D49**, 1378 (1994); W. A. Ponce, A. Zepeda and R. G. Lozano, Phys. Rev. **D49**, 4954 (1994).
- [9] Y. Koide, Phys. Rev. **D60**, 077301 (1999).
- [10] Y. Koide, Phys. Rev. **D73**, 057901 (2006).

- [11] C. Froggatt and H. B. Nielsen, Nucl. Phys. **B147**, 277 (1979).
- [12] S. Pakvasa and H. Sugawara, Phys. Lett. **B73**, 61 (1978); H. Harari, H. Haut and J. Weyers, Phys. Lett. **B78**, 459 (1978); E. Derman, Phys. Rev. **D19**, 317 (1979); D. Wyler, Phys. Rev. **D19**, 330 (1979).
- [13] Y. Koide, e-print arXiv: hep-ph/0005137 (unpublished).
- [14] S. N. Ahmed *et al.*, SNO Collaboration, Phys. Rev. Lett. **92**, 181301 (2004); T. Araki *et al.* KamLAND Collaboration, Phys. Rev. Lett. **94**, 081801 (2005); A. W. P. Poon, in Proceedings of the XXII International Symposium on Lepton and Photon Interactions at High Energies, Uppsala, Sweden, 30 June – 5 July, 2005, edited by R. Brenner, C. P. Heros and J. Rathsmann (World Scientific Pub. 2006), p. 278.
- [15] J. Nelson, MINOS collaboration, presented in Neutrino 2006, at Santa Fe. Also see, J. Hosaka *et al.* the Super-Kamiokande Collaboration, arXiv: hep-ex/0604011.
- [16] Y. Koide, Phys. Rev. **D64**, 053014 (2001).
- [17] Y. Koide, Eur. Phys. J. **C9**, 335 (1999); Phys. Rev. **D61**, 035008 (2000).
- [18] P. F. Harrison, D. H. Perkins and W. G. Scott, Phys. Lett. **B458**, 79 (1999); Phys. Lett. **B530**, 167 (2002); Z. Z. Xing, Phys. Lett. **B533**, 85 (2002); P. F. Harrison and W. G. Scott, Phys. Lett. **B535**, 163 (2003); Phys. Lett. **B557**, 76 (2003); E. Ma, Phys. Rev. Lett. **90**, 221802 (2003); C. I. Low and R. R. Volkas, Phys. Rev. **D68**, 033007 (2003); X.-G. He and A. Zee, Phys. Lett. **B560**, 87 (2003).
- [19] T. Fukuyama and H. Nishiura, hep-ph/9702253, in Proceedings of *the International Workshop on Masses and Mixings of Quarks and Leptons*, Shizuoka, Japan, 1997, edited by Y. Koide (World Scientific, Singapore, 1998), p. 252; E. Ma and M. Raidal, Phys. Rev. Lett. **87** (2001) 011802; C. S. Lam, Phys. Lett. **B507** (2001) 214; K. R. S. Balaji, W. Grimus and T. Schwetz, Phys. Lett. **B508** (2001) 301; W. Grimus and L. Lavoura, Acta Phys. Pol. **B32** (2001) 3719.
- [20] E. Ma, Phys. Rev. **D73**, 057304 (2006).
- [21] H. Fusaoka and Y. Koide, Phys. Rev. **D57**, 3986 (1998).

Table 1: Values of $R_f(\eta_1, \eta_2)$ which is defined by Eq. (4.2): For convenience, the pole mass values [1] for the charged lepton masses, the values (3.13) for the neutrino masses, and the quark mass values for the running mass values at $\mu = m_Z$ [21] are used, respectively. In the Table, the center values have been used as the input values.

Sector	$R(+, +)$	$R(-, +)$	$R(+, -)$	$R(-, -)$
Charged lepton	0.999998	0.946922	0.376006	0.34374
Neutrino	1.28	1.00	0.251	0.137
Up-quark	0.753	0.743	0.590	0.581
Down-quark	0.955	0.834	0.481	0.397




Fig. 1 R versus θ_ν under the relation (1.2). The horizontal lines denote the observed values $R = (2.9 \pm 0.5) \times 10^{-2}$.